



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Br

Search PubMed for Go Clear

Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

Entrez PubMed  
Overview  
Help | FAQ  
Tutorial  
New/Noteworthy  
E-Utilities


PubMed Services  
Journals Database  
MeSH Database  
Single Citation Matcher  
Batch Citation Matcher  
Clinical Queries  
LinkOut  
Cubby

Related Resources  
Order Documents  
NLM Gateway  
TOXNET  
Consumer Health  
Clinical Alerts  
ClinicalTrials.gov  
PubMed Central

Privacy Policy

☐ 1: Expert Rev Anticancer Ther. 2004 Feb;4(1):18-26.

Related Articles, Link

**Iodine-131 tositumomab (Bexxar(R)): radioimmunoconjugate therapy for indolent and transformed B-cell non-Hodgkin's lymphoma.****Friedberg JW, Fisher RI.**

Lymphoma Program, James P Wilmot Cancer Center, University of Rochester, Rochester, NY, USA. richard\_fisher@urmc.rochester.edu

Tositumomab is an immunoglobulin G murine monoclonal antibody that binds to the CD20 antigen on the surface of normal and malignant human B-cells. Tositumomab is linked covalently with iodine-131 to produce the radioimmunoconjugate iodine-131 tositumomab (Bexxar(R)). The iodine-131 tositumomab regimen was approved by the US Food and Drug Administration in June 2003 for the treatment of patients with CD20-positive follicular non-Hodgkin's lymphoma, both with and without transformation, whose disease is refractory to rituximab (Rituxan(R)) and has relapsed following chemotherapy. The dose-limiting toxicity of iodine-131 tositumomab is bone marrow suppression and resulting cytopenias. Unlike chemotherapy, the majority of nonhematologic adverse events associated with iodine-131 tositumomab are mild to moderate in nature and usually self limited. Iodine-131 tositumomab represents one of the most active single agents for the treatment of recurrent indolent and transformed B-cell non-Hodgkin's lymphoma, as demonstrated by several clinical trials summarized in this review. At the present time, the use of radioimmunoconjugate therapy is largely limited to patients with disease refractory to rituximab therapy and transformed disease not amenable to high-dose therapy and autologous stem cell support. Longer follow-up of ongoing clinical trials should provide reassurance as to safety and insights as to the additive stem cell toxicity from iodine-131 tositumomab administration. Studies are also addressing the role of iodine-131 tositumomab as a component of initial therapy for indolent non-Hodgkin's lymphoma and in additional histologies of non-Hodgkin's lymphoma.

PMID: 14748653 [PubMed - in process]

20F2

09/26/2, 587

[89] The BEXXAR<sup>®</sup> Therapeutic Regimen (Tositumomab and Iodine I 131 Tositumomab) Produced Durable Complete Remissions in Heavily Pretreated Patients with Non-Hodgkin's Lymphoma (NHL), Rituximab-Relapsed/Refractory Disease, and Rituximab-Naïve Disease. Session Type: Oral Session

Morton Coleman, Mark S. Kaminski, Susan J. Knox, Andrew D. Zelenetz, Julie M. Vose (Intr. by Morton Coleman) Weill Medical College of Cornell University, New York, NY, USA; University of Michigan Cancer Center, Ann Arbor, MI, USA; Stanford University Medical Center, Stanford, CA, USA; Memorial Sloan-Kettering Cancer Center, New York, NY, USA; University of Nebraska Medical Center, Omaha, NE, USA

**Introduction:** The BEXXAR therapeutic regimen was recently approved for the treatment of patients with CD20-positive, follicular NHL, with and without transformation, whose disease is refractory to rituximab and who have relapsed following chemotherapy. Approval was based on a study that included 40 patients, 35 (88%) of whom met the definition of rituximab-refractory (no response or a duration of response of  $\leq 6$  months). The response rate for the entire population was 68%, with a median duration of response of 16 months. The results of this study were supported by demonstration of durable objective responses in 4 other studies enrolling 190 rituximab-naïve patients with follicular NHL, with or without transformation, who were relapsed/refractory (rel/ref) to chemotherapy. The response rates in these 4 studies ranged from 47% to 64%, with median durations of response ranging from 12 to 18 months. **Methods:** The goal of the study was to determine the number of patients who achieved long-term durable complete responses (CR) in both the rituximab-rel/ref group and the rituximab-naïve group and to assess their durations of response. A durable CR was defined as a CR with a progression-free survival (PFS) of  $\geq 12$  months as assessed by a blinded, independent panel of oncologists and radiologists (MIRROR Panel). **Results:** Of the 230 patients evaluable for response, 55 (24%) met the definition of a durable CR (11 of 40 [28%] from the rituximab-rel/ref study, and 44 of 190 [23%] from the 4 studies in rituximab-naïve patients).

#### Patient Demographics for Durable Complete Responders

	Rituximab naïve patients	Rituximab rel/ref patients
Median age years, (range)	52 (23-82)	53 (39-73)
Stage III/IV at study entry, %	93	73
Transformed histology, %	27	9
Mean prior therapies, n	3.1	4.1
Prior radiotherapy, %	18	27
Maximum tumor diameter $\geq 5$ cm, %	44	18
Median follow-up, years (range)	5.0 (1.3-9.5)	3.9 (3.0-4.6)

Among rituximab-naïve patients, median duration of response was 4.9 years. Among rituximab-rel/ref patients, median duration of response has not been reached, with a median follow-up of 3.9 years. Thirty-three of 44 (75%) patients with durable CRs in the rituximab-naïve population continue in CR at their last assessment. Eight of 11 (73%) patients with durable CRs in the rituximab-rel/ref group continue in CR at their last assessment. **Conclusions:** Independently assessed durable CRs were noted with similar frequency in patients with rituximab-rel/ref disease (28%) and rituximab-naïve patients with chemotherapy-rel/ref disease (23%). With a median follow-up of 4.6 years, 75% of patients with durable CRs continue in complete remission. The BEXXAR therapeutic regimen produced durable CRs in patients with follicular NHL who had relapsed following rituximab or who were refractory to rituximab, and in patients who had never received rituximab. Abstract #89 appears in Blood, Volume 102, issue 11, November 16, 2003

**Keywords:** rituxan failure|first-line therapy|BEXXAR

Sunday, December 7, 2003 4:15 PM

Simultaneous Session: Chemotherapy/Rituximab Combinations and Radiolimmunotherapy (4:15 PM-5:45 PM)

Close Window

1 of 2

## Treatment With Ibritumomab Tiuxetan Radioimmunotherapy in Patients With Rituximab- Refractory Follicular Non-Hodgkin's Lymphoma

By Thomas E. Witzig, Ign W. Flinn, Leo I. Gordon, Christos Emmanouilides, Myron S. Czuczman, Mansoor N. Saleh, Larry Cripe, Gregory Wiseman, Teresa Olejnik, Pratik S. Multani, and Christine A. White

**Purpose:** Rituximab is commonly used as a single agent or in combination therapy for non-Hodgkin's lymphoma (NHL). Ibritumomab tiuxetan radioimmunotherapy targets the same antigen as rituximab and has demonstrated efficacy in rituximab-naïve NHL. This study evaluated ibritumomab tiuxetan in the treatment of rituximab-refractory follicular NHL.

**Patients and Methods:** Eligible patients were refractory to rituximab; this was defined as no objective response to rituximab (375 mg/m<sup>2</sup> weekly for 4 weeks) or time to progression (TTP) of  $\leq$  6 months. The ibritumomab tiuxetan treatment regimen consisted of pretreatment with rituximab (250 mg/m<sup>2</sup> intravenously on days 1 and 8) to deplete peripheral blood B cells, then yttrium-90 ibritumomab tiuxetan (0.4 mCi/kg; maximum, 32 mCi) intravenously on day 8, administered on an outpatient basis. An imaging/dosimetry dose of indium-111 ibritumomab tiuxetan (5 mCi) was injected after rituximab (day 1) in 28 patients.

**Results:** Fifty-seven patients were treated. The median age was 54 years, 74% had tumors  $\geq$  5 cm, and all were extensively pretreated (median, four prior therapies; range, one to nine). The estimated radiation-absorbed doses to healthy organs were below the study-defined limit in all patients studied with dosimetry. The overall response rate for the 54 patients with follicular NHL was 74% (15% complete responses and 59% partial responses). The Kaplan-Meier-estimated TTP was 6.8 months (range, 1.1 to  $\geq$  25.9 months) for all patients and 8.7 months for responders. Adverse events were primarily hematologic; the incidence of grade 4 neutropenia, thrombocytopenia, and anemia was 35%, 9%, and 4%, respectively.

**Conclusion:** Ibritumomab tiuxetan radioimmunotherapy is effective in rituximab-refractory patients. The only significant toxicity is hematologic.

*J Clin Oncol* 20:3262-3269. © 2002 by American Society of Clinical Oncology.

**R**ITUXIMAB (RITUXAN; IDEC Pharmaceuticals, San Diego, CA, and Genentech, Inc, South San Francisco, CA) is a chimeric anti-CD20 monoclonal antibody that was approved in 1997 by the United States Food and Drug Administration for the treatment of relapsed or refractory low-grade or follicular non-Hodgkin's lymphoma (NHL). A 48% overall response rate (ORR) and a time to progression (TTP) of approximately 1 year were demonstrated in the pivotal trial of rituximab.<sup>1</sup> Recent studies have indicated that responses can be reinduced on retreatment with rituximab<sup>2</sup> and that rituximab can be used concomitantly with chemotherapy.<sup>3-6</sup>

Technical advances have made it possible to link radionuclides such as yttrium-90 (<sup>90</sup>Y) to monoclonal antibodies

specifically to target radiation to lymphoma cells. Yttrium-90 is a beta-emitting radionuclide that delivers 90% of its radiation (2.3 MeV) over a mean path length of 5 mm and has a half-life of 64 hours. These characteristics are particularly advantageous for treating bulky, poorly vascularized tumors and those with heterogeneous antigen expression.

Ibritumomab tiuxetan (Zevalin; IDEC Pharmaceuticals) is a short-course radioimmunotherapy that uses immunobiologic and radiolytic mechanisms of action to destroy both dividing and nondividing tumor cells. Ibritumomab is the murine, parent anti-CD20 antibody that was engineered to develop rituximab. Tiuxetan is a linker/chelator covalently attached to ibritumomab. Ibritumomab tiuxetan can chelate indium-111 (<sup>111</sup>In) for imaging or <sup>90</sup>Y for therapy. Thus, the antibody specifically targets radiation to CD20<sup>+</sup> cells while sparing normal nonlymphoid cells.

Because rituximab immunotherapy is widely used to treat patients with NHL, a critical issue is whether patients treated previously with rituximab can respond to ibritumomab tiuxetan. This study tested the hypothesis that tumor resistance to an unconjugated anti-CD20 antibody alone can be overcome by subsequent CD20-directed radioimmunotherapy.

### PATIENTS AND METHODS

#### Eligibility

Eligible patients with follicular B-cell NHL had prior treatment with rituximab, 375 mg/m<sup>2</sup> once weekly for 4 weeks, and either did not

From the Mayo Clinic, Rochester, MN; Johns Hopkins Oncology Center, Baltimore, MD; Northwestern University/Robert H. Lurie Cancer Center, Chicago, IL; University of California, Los Angeles, and IDEC Pharmaceuticals Corporation, San Diego, CA; Roswell Park Cancer Institute, Buffalo, NY; University of Alabama, Birmingham, AL; and Indiana Cancer Pavilion, Indianapolis, IN.

Submitted November 5, 2001; accepted April 10, 2002.

Supported by IDEC Pharmaceuticals Corp, San Diego, CA.

Address reprint requests to Thomas E. Witzig, MD, 620 Stabile Bldg, Mayo Clinic, Rochester, MN 55905; email: witzig@mayo.edu.

© 2002 by American Society of Clinical Oncology.

0732-183X/02/2015-3262/\$20.00

respond or had a TTP of less than 6 months. The follicular NHL histologic subtype (Revised European-American Lymphoma classification,<sup>7</sup> follicle center grade 1, 2, or 3) was confirmed by biopsy. The study prospectively allowed enrollment of patients with small lymphocytic or transformed NHL who were rituximab nonresponders from the control arm of a randomized ibritumomab tiuxetan trial,<sup>8</sup> although such patients were to be included in the analysis of safety only.

Patients were to be at least 18 years old and have bidimensionally measurable disease with at least one lesion measuring  $\geq 2.0$  cm in a single dimension, less than 25% bone marrow involvement with lymphoma. World Health Organization performance status  $\leq 2$ , absolute neutrophil count (ANC)  $\geq 1,500$  cells/mm<sup>3</sup>, platelet count  $\geq 150,000$  cells/mm<sup>3</sup>, serum creatinine and total bilirubin  $\leq 2$  mg/dL each, expected survival  $\geq 3$  months, no antineoplastic therapy for 3 weeks (6 weeks if treated with nitrosourea or mitomycin), and no hematopoietic growth factor treatment (granulocyte colony-stimulating factor or granulocyte-macrophage colony-stimulating factor) within 2 weeks of study entry. Patients with prior autologous bone marrow transplantation or peripheral blood stem-cell support, prior radioimmunotherapy, or external-beam radiation therapy to more than 25% of active bone marrow were not eligible. Also, patients were to have no measurable serum human antimurine or antichimeric antibodies (HAMA or HACA, respectively). Patients were required to sign written, informed consent, and each clinical site had to obtain institutional review board approval before conducting this study.

#### Study Design

Patients were to receive one course of standard ibritumomab tiuxetan outpatient treatment, as described previously<sup>9</sup>; no hospitalization, patient isolation, or shielding was required. Briefly, this consisted of an infusion of rituximab 250 mg/m<sup>2</sup> on day 1 and day 8 (range, days 7 to 9) and <sup>90</sup>Y ibritumomab tiuxetan (0.4 mCi/kg [15 MBq/kg]; maximum, 32 mCi [1.2 GBq]) on day 8 (range, days 7 to 9) immediately after the second rituximab infusion. Radioincorporation of <sup>90</sup>Y or <sup>111</sup>In into the immunoconjugate was to be performed at each clinical site or at an affiliated commercial radiopharmacy. Acceptable radioincorporation was to be  $\geq 95\%$  for both <sup>90</sup>Y and <sup>111</sup>In radionuclides, as determined with an instant thin-layer chromatographic method.<sup>10,11</sup>

Radiation dosimetry, performed as defined prospectively in the protocol, was to include an imaging/dosimetry dose of <sup>111</sup>In ibritumomab tiuxetan (5 mCi; 185 MBq) to be administered immediately after the rituximab infusion on day 1. Dosimetry was to be performed at the clinical site to ensure that no patient received an estimated radiation-absorbed dose greater than 20 Gy to healthy organs and greater than 3 Gy to red marrow. The protocol was amended during enrollment to remove the dosimetry requirement.

For the purpose of collecting safety data, a treatment period was defined as the time from the rituximab infusion on day 1 to 12 weeks after <sup>90</sup>Y ibritumomab tiuxetan. Disease status evaluations included a combination of medical history, physical examination, bone marrow biopsy, and computed tomography scan or magnetic resonance imaging of the neck, chest, abdomen, and pelvis, as well as other clinically relevant information. Before treatment, lesions were measured bidimensionally, and the sum of the products of the longest perpendicular diameters (SPD) was calculated as a reference to determine baseline disease status. Posttreatment follow-up evaluations were to be conducted at 1 month and then at 3-month intervals for the first 2 years, and at 6-month intervals for the next 2 years or until disease progression. Safety follow-up continued until disease progression necessitated intervention with another antineoplastic therapy. Efficacy (intent to treat) was to be

evaluated in patients with follicular histology only, whereas safety analyses (intent to treat) were to include all enrolled patients.

Patients were evaluated for response by an independent, third-party panel of radiologists and oncologists who were expert in lymphoma (Lymphoma Experts Confirmation of Response). The panel was blinded to the investigator's response assessment. Response was classified according to protocol-defined criteria and International Workshop NHL response criteria.<sup>12</sup> Response is reported here by using International Workshop NHL response criteria, because those are now the accepted standards.

#### Statistical Methods

The primary efficacy end point was a target ORR (complete response [CR] plus continuous complete remission plus partial response [PR]) of at least 35% in patients with follicular NHL as assessed by Lymphoma Experts Confirmation of Response. Secondary efficacy end points included TTP and duration of response (DR). TTP was defined as the time from the date of the first infusion given on day 1 in this study to the date of disease progression; DR was calculated from the date of the first response observation to the date of disease progression. Time-to-event variables were estimated with the Kaplan-Meier product-limits method.<sup>13</sup>

An additional secondary efficacy end point prospectively defined in the protocol was the comparison of the ORR and DR with the ORR and DR obtained with the prior rituximab therapy and with the last prior chemotherapy, by using McNemar's test. A second exploratory analysis comparing ibritumomab tiuxetan with prior chemotherapy and prior rituximab therapy by using the sign test and signed-rank test was performed at the request of the Food and Drug Administration. The analysis was defined in the following manner: (1) ibritumomab tiuxetan was favored if a patient responded to ibritumomab tiuxetan but not to rituximab (or chemotherapy), and if a patient responded to both, the DR to ibritumomab tiuxetan must have been  $\geq 3$  months longer than the response to the prior therapy; (2) rituximab (or chemotherapy) was favored if a patient responded to rituximab (or chemotherapy) but not to ibritumomab tiuxetan, and if a patient responded to both, the DR to rituximab (or chemotherapy) must have been  $\geq 3$  months longer than the response to ibritumomab tiuxetan; and (3) the response was classified as neutral if a patient did not respond to either therapy or responded to both but the DR to ibritumomab tiuxetan was within 3 months of the DR to both.

The patients' perceived quality of life was assessed with a self-rating instrument, the Functional Assessment of Cancer Therapy-General questionnaire,<sup>14</sup> at baseline and at 12 weeks after treatment. Differences in the total score were evaluated with the paired *t* test.

Safety assessments were based on analysis of adverse events and clinical laboratory data by using the National Cancer Institute adult toxicity criteria, version 2.0. Laboratory assessments included hematology and blood chemistry data, serum HAMA/HACA, and monitoring of peripheral blood lymphocyte subpopulations with flow cytometry (B cells were defined as those expressing the CD19 cell-surface antigen). The duration of hematologic toxicity for each hematologic variable (ANC, platelet count, and hemoglobin concentration) was calculated by using two methods. In method A, duration was measured from the date of the last laboratory value before the development of grade 3 or 4 toxicity to the date of the next value in grade 2 after the nadir. In method B, duration was measured from the date of the first laboratory value in grade 3 or 4 toxicity to the date of the last value in grade 3 after the nadir. Baseline residual serum rituximab concentrations were measured before the rituximab infusion on day 1.

An exploratory analysis was conducted to determine whether a relationship existed between a patient's baseline bone marrow involve-

ment with NHL as determined by light microscopy and grade 4 hematologic toxicity after treatment with ibritumomab tiuxetan. Patients were sorted into four groups on the basis of bone marrow involvement: 0% bone marrow involvement, 0.1% to 5.0% bone marrow involvement, 5.1% to less than 20% bone marrow involvement, and  $\geq 20\%$  bone marrow involvement. *P* values were generated by Fisher's exact two-tailed test.

## RESULTS

### Patient Characteristics

Fifty-seven patients were enrolled and completed treatment at 18 clinical sites between July 7, 1998, and October 12, 1999. Patient characteristics are listed in Table 1. Fifty-four patients had follicular NHL (grade 1, 28 patients [52%]; grade 2, 21 patients [39%]; and grade 3, five patients [9%]), two patients had small lymphocytic NHL, and one had transformed diffuse large-cell NHL. Patients were extensively pretreated, and 74% had tumors  $\geq 5$  cm.

### Dosimetry

<sup>111</sup>In ibritumomab tiuxetan imaging and dosimetry were performed in 28 patients, and the complete results were reported separately.<sup>15</sup> The estimated radiation-absorbed doses were within acceptable levels, allowable for proceeding with <sup>90</sup>Y ibritumomab tiuxetan treatment in all patients. The median estimated radiation-absorbed doses were 8.1 Gy to the spleen (range, 4.2 to 23.0 Gy); 5.1 Gy to the liver (range, 2.6 to 12.0 Gy); 2.0 Gy to the lungs (range, 1.4 to 5.3 Gy); 0.22 Gy to the kidneys (range,  $< 0.01$  to 0.66 Gy); and 0.74 Gy to red marrow (range, 0.29 to 1.2 Gy).

### Efficacy

The ORR was 74% (40 of 54 patients), with 15% (eight of 54) CRs and 59% (32 of 54) PRs. Ibritumomab tiuxetan produced a tumor shrinkage (reduction in the SPD) in 94% (51 of 54) of patients. The mean reduction in SPD was 74% in the 32 patients who achieved a PR and was 34% in the 14 patients with stable disease.

The ORR to ibritumomab tiuxetan therapy was not statistically different ( $P < .05$ ) in patients with or without bone marrow involvement (81% v 71%), in those with or without splenomegaly (83% v 73%), and in patients weighing  $\leq 80$  kg who were dosed per kilogram or patients weighing more than 80 kg who received a capped dose of 32 mCi (77% v 68%). Half of patients with bulky disease more than 10 cm and 68% who received four or more prior antineoplastic regimens responded to ibritumomab tiuxetan. No statistically significant difference was found between the response of patients with measurable serum rituximab concentrations (66%; 25 of 38) and those with undetectable serum rituximab concentrations (93%; 13 of 14 [ $P = .078$ ]).

Table 1. Patient Characteristics (N = 57)

Characteristic	No.	%
Lymphoma histology		
Follicular	54	95
Small lymphocytic	2	4
Transformed large-cell	1	2
Age, years		
Median	54	
Range	34-73	
Female	29	51
White	54	95
Disease stage at study entry		
I/II	6	10
III/IV	51	90
Tumor size		
$< 5$ cm	15	26
5 to $< 7$ cm	17	30
7 to $< 10$ cm	14	25
$\geq 10$ cm	11	19
Splenomegaly	7	12
Bone marrow involvement	18	32
Two or more extranodal disease sites	10	18
WHO performance status		
0, 1	54	95
2	3	5
IPi risk group (if known)		
Low	25	44
Low/intermediate	12	21
Intermediate/high	7	12
High	4	7
B-cell count in peripheral blood		
Undetectable	21	37
Low ( $< 32$ cells/mm <sup>3</sup> )	20	35
Normal/high ( $\geq 32$ cells/mm <sup>3</sup> )	14	25
Unknown	2	4
Median no. of prior antineoplastic therapy regimens	4	
Resistance to chemotherapy*	1-9	
Last therapy	37	67
Any therapy	45	82
Prior serum rituximab concentration†		
Undetectable	15	26
1 to $< 10$ $\mu$ g/mL	30	53
$\geq 10$ $\mu$ g/mL	9	16
Unknown	3	5

Abbreviations: WHO, World Health Organization; IPi, International Prognostic Index.

\*Resistance: nonresponders or patients who had disease progression within 6 months.

†Measured before rituximab infusion on day 1.

although a trend was demonstrated. Peripheral blood B cells were quantitated at baseline by using flow cytometry in all 54 patients with follicular NHL. Forty-seven percent (nine of 19 patients) with no detectable B cells responded to ibritumomab tiuxetan compared with 88% (29 of 33) with measurable B cells ( $P = .003$ ).

# RADIOIMMUNOTHERAPY FOR RITUXIMAB-REFRACTORY NHL

3265




Fig 1. Kaplan-Meier analysis of time to progression for patients with follicular histology: ( $\Delta$ ) intent-to-treat patients,  $n = 54$ ; (+) responders,  $n = 40$ . C, censored (30%).

The median TTP estimated by the Kaplan-Meier method was 6.8 months (range, 1.1 to  $\geq 25.9$  months) with 30% of the data censored (Fig 1). The median TTP in the 40 responders was 8.7 months (range, 1.7 to  $\geq 25.9$  months), with 28% of the data censored. The median DR estimated by the Kaplan-Meier method was 6.4 months (range, 0.5 to  $\geq 24.9$  months).

Figure 2 displays the ORR achieved with ibritumomab tiuxetan compared with that achieved with prior rituximab and the last chemotherapy. Of the 17 patients who responded to prior rituximab treatment, 88% (15 of 17) responded to ibritumomab tiuxetan, and of the 34 patients who responded to the last chemotherapy, 82% (28 of 34) responded to ibritumomab tiuxetan. In the patients who




Fig 2. (A) Ibritumomab tiuxetan was the median fifth regimen; chemotherapy was the median third-regimen treatment. (B) Comparison of ibritumomab tiuxetan with last prior chemotherapy and prior rituximab.

responded to prior rituximab treatment, the median DR was 11.5 months for ibritumomab tiuxetan (46% of patients censored) and 3.0 months for rituximab (no patient censored;  $P = .001$ ). In the 25 patients in whom DR information was available for the last chemotherapy, the median DR was 5.4 months for ibritumomab tiuxetan (16% of patients censored) and 5.0 months for last chemotherapy (no patient censored;  $P = .76$ ). Table 2 displays this comparison based on response to treatment and DR in a combined analysis as described in the Patients and Methods, under Statistical Methods.

**Functional Assessment of Cancer Therapy**—General quality-of-life survey data were available for 20 patients. The mean score of 85 before ibritumomab tiuxetan treatment improved significantly to 92.8 at 12 weeks after treatment ( $P = .003$ ). Disease-related symptoms improved or resolved in all responders.

## Safety

Adverse events after ibritumomab tiuxetan treatment were primarily hematologic and transient; no patient discontinued treatment because of an adverse event. The duration of hematologic toxicity is listed in Table 3.

The incidence of grade 4 neutropenia, thrombocytopenia, and anemia was 35%, 9%, and 4%, respectively. With Fisher's exact two-tailed test, as described under Statistical Methods, the incidence of the grade 4 ANC nadir correlated statistically with increasing bone marrow involvement with NHL ( $P = .004$ ). In particular, 28% of patients with no bone marrow involvement developed grade 4 ANC nadirs, whereas 100% of patients with 20% to 25% bone marrow involvement developed grade 4 ANC nadirs. No statistically significant difference in grade 4 hematologic toxicity was observed between patients weighing  $\leq 80$  kg and patients weighing more than 80 kg ( $P = .526$  to .999 for hemoglobin, ANC, and platelets). No correlation existed between hematologic nadirs and either baseline serum rituximab concentrations or baseline peripheral blood B-cell counts. Eighteen patients (32%) received growth factor therapy; 11 (19%) received erythropoietin, five (9%) received granulocyte colony-stimulating factors, and one (2%) received a platelet growth factor. Thirteen patients (23%) received platelet transfusions, and 14 (25%) received RBC transfusions.

The most common nonhematologic events during the treatment period were infusion related (Table 4) and were consistent with those described for rituximab infusion.<sup>1</sup> These included asthenia in 54% (31 of 57 patients), nausea in 35% (20 of 57), chills in 25% (14 of 57), and fever in 21% (12 of 57). Eighty-three adverse events occurred on a treatment day (66 on day 1 and 22 on day 8); all were grade 1 or 2 except one (grade 3 tumor pain). Four patients (7%) were hospitalized, one each with febrile neutropenia, pneu-

Table 2. Additional Comparison of Ibritumomab Tiuxetan Versus Prior Rituximab Therapy or Last Chemotherapy (n = 54)\*

Variable	Favors, † (%)		Neutral (%)	P	
	Ibritumomab Tiuxetan	Prior Therapy		Signed-Rank Test	Sign Test
Ibritumomab tiuxetan versus prior rituximab	48	9	43	< .001	.011
Ibritumomab tiuxetan versus last chemotherapy*	30	30	41	.388	.377

\*Ibritumomab tiuxetan was the median fifth regimen; chemotherapy was the median third-regimen treatment.

†Defined as (1) response to one therapy but not to the other, or if response to both, the DR must be  $\geq 3$  months longer than response to the other therapy; (2) neutral if a patient did not respond to either therapy or responded to both but the DR to ibritumomab tiuxetan was within 3 months of the DR to both (see Patients and Methods).

monia/sepsis, cellulitis, and urinary tract infection. No patient developed HACA, and one patient developed a low-titer HAMA (29  $\mu\text{g/mL}$ ) on day 42. This patient experienced no adverse events attributable to HAMA and achieved a PR. One patient with a detectable HACA titer at study entry (44.7  $\text{ng/mL}$ ) received an exemption to allow treatment. This patient did not respond to ibritumomab tiuxetan and experienced no grade 3 or 4 or other serious adverse event.

One patient died as a result of acute myelogenous leukemia (AML) approximately 14 months after treatment. The patient had previously received single-agent and combination chemotherapy, including chlorambucil; cyclophosphamide, doxorubicin, vincristine, and prednisone; and prednisone, methotrexate, doxorubicin, cyclophosphamide, and etoposide/mechlorethamine, vincristine, procarbazine, and prednisone. Blasts were detected in the patient's peripheral blood during the first month after ibritumomab tiuxetan treatment. AML was diagnosed at month 8, when cytogenetic studies demonstrated monosomy 7, inversion 3q, and t(21;22). Subsequent analysis of pretreatment bone marrow for this patient revealed 23% of cells with monosomy 7; however, this was within the normal range for the assay (upper normal limit, 25%). One patient taking oral anticoagulant therapy for a history of deep vein thrombophlebitis and self-prescribed ibuprofen died on study day 71 as a result of a traumatic subdural hematoma that occurred at platelet nadir.

## DISCUSSION

Patients with advanced-stage follicular NHL are likely to die as a result of that disease.<sup>16</sup> Most patients respond to subsequent treatments; however, recurrence, progression, transformation to a more aggressive histology, or a combination of these is typically unavoidable.<sup>17</sup> Patients usually have repeated relapses, and subsequent courses of chemotherapy characteristically lead to fewer and shorter remissions.<sup>18</sup> Cumulative toxicity associated with cytotoxic chemotherapy often necessitates dose reduction, shorter courses, or drug substitutions, each of which may compromise response to treatment. Rituximab immunotherapy has been an important treatment advance in NHL because of its efficacy, short duration of therapy, and acceptable toxicity profile. The aim of this study was to evaluate whether rituximab-refractory patients could achieve responses with an anti-CD20 radioimmunotherapy directed against the same epitope as rituximab.

Fifty-four patients with rituximab-refractory follicular NHL received the ibritumomab tiuxetan regimen and achieved a 74% ORR (15% CR). This response rate exceeded the protocol-targeted 35% ORR and is particularly encouraging in this patient population, which was extensively pretreated (median of four regimens) with chemotherapy. By using two separate analysis methods, ibritumomab tiuxetan efficacy was superior to prior rituximab therapy and compared favorably with last chemotherapy.

Table 3. Duration of Hematologic Toxicity

Variable	Baseline	Nadir	Days From Baseline to Nadir	Median Duration for Patients With Grade 3 or 4 Nadir (days)*	
				Method A	Method B
ANC, cells/mm <sup>3</sup>	3,500	700	63	22	8
Platelets, cells/mm <sup>3</sup>	214,000	33,000	55	24	12
Hemoglobin, g/dL	13.3	9.9	68	8	1

\*ANC  $< 1,000$  cells/mm<sup>3</sup>; platelets  $< 50,000$  cells/mm<sup>3</sup>; and hemoglobin  $< 8$  g/dL. Method A, duration measured from the last date in grade 2 before nadir to the first date in grade 2 after nadir; method B, duration measured from the first date in grade 3 or 4 before nadir to the last date in grade 3 or 4 after nadir.

Table 4. Incidence of Most Common and Grade 3 or 4 Nonhematologic Adverse Events (N = 57)\*

Event	Grade				Total	
	1	2	3	4	No.	%
Body as a whole						
Asthenia	17	12	2	0	31	54
Chills	12	2	0	0	14	25
Fever	5	6	1	0	12	21
Headache	4	3	0	0	7	12
Throat irritation	6	1	0	0	7	12
Abdominal pain	3	4	0	0	7	12
Neck pain	1	0	1	0	2	4
Tumor pain	0	0	1	0	1	2
Sepsis	0	0	0	1	1	2
Cardiovascular system						
Deep vein thrombophlebitis	0	0	1	0	1	2
Digestive system						
Nausea	17	3	0	0	20	35
Pancytopenia	0	0	0	1	1	2
Respiratory system						
Dyspnea	6	1	0	0	7	12
Pneumonia	0	0	0	1	1	2
Urogenital system						
Vaginal hemorrhage	0	0	1	0	1	2

\*The incidence was  $\geq 10\%$  of patients and all grade 3 or 4 adverse events that occurred during the treatment period and classified as probably or possibly related to study drug or relationship unknown. Each patient was counted only once under the worst-grade adverse event experienced, excluding neutropenia, leukopenia, thrombocytopenia, and anemia.

The myelosuppression associated with ibritumomab tiuxetan is gradual in onset and occurs later than typical chemotherapy-induced myelosuppression. Despite myelosuppression, infections were unusual; only 7% of patients were hospitalized, and there were no infection-related deaths. The correlation between percentage of bone marrow involvement with lymphoma and hematologic toxicity confirmed that hematologic toxicity was related to specific targeting of  $^{90}\text{Y}$  ibritumomab tiuxetan to marrow and secondary irradiation of marrow hematopoietic cells. The majority of nonhematologic events consisted of grade 1 or 2 constitutional symptoms such as asthenia, nausea, fever, and chills associated with infusion days. Similarly, the most common grade 3 or 4 nonhematologic events were asthenia and pain and were experienced by only 11% of patients.

One patient developed AML and died approximately 14 months after ibritumomab tiuxetan treatment. Baseline marrow cytogenetic data revealed possible pretreatment chromosomal abnormalities, and blasts were apparent in the patient's peripheral blood as early as the first month of the study, suggesting that the condition was incipient. Secondary AML has been observed with increasing frequency among NHL patients, and the contribution of prior alkylator therapy seems unequivocal.<sup>19-23</sup> Moreover, the possibility that certain NHL patients are prone to the development of secondary AML has not been ruled out.<sup>19</sup> It will be

important to observe patients carefully for the development of late marrow toxicity.

A primary concern of re-treatment with monoclonal antibody-based therapy is the development of antixenogenic antibodies (HAMA and HACA) that may reduce the therapeutic antibody serum half-life and increase the risk of adverse events such as fever, hypotension, rash, arthritis and arthralgias, and nerve palsies.<sup>24-26</sup> However, only one patient (< 2%) developed HAMA in this study, and this patient achieved a PR. This low incidence of HAMA and HACA responses is consistent with findings in other ibritumomab tiuxetan trials.<sup>9,27-29</sup> One additional patient with a HACA titer at baseline experienced no unusual toxicity or adverse events after treatment with ibritumomab tiuxetan.

Some patients had received rituximab shortly before entering this study. Prior treatment with rituximab did not compromise patient safety; all normal organ doses were acceptable and were consistent with those observed in other ibritumomab tiuxetan studies.<sup>30,31</sup> The presence of residual serum rituximab or the absence of peripheral blood B cells at baseline did not preclude tumor response, although response rates tended to be lower for patients with these factors. As demonstrated previously for both rituximab-refractory and rituximab-naïve patients, undetectable B cells at baseline predicted a shorter disease-free interval and poor prognosis.<sup>32</sup> However, no measurable difference in



hematologic toxicity was noted in patients stratified by either residual serum rituximab concentration or baseline peripheral blood B-cell count as measured by grade 4 ANC, platelet, and hemoglobin nadir values. Therefore, it seems that the trend toward lower response rates is not due to altered biodistribution, but that these unfavorable baseline factors select for a poorer-prognosis population. Patients with these factors may nevertheless achieve a favorable tumor response.

This study demonstrates that patients can be safely and effectively treated with ibritumomab tiuxetan after prior treatment with rituximab. The challenge in future studies will be to evaluate whether high-dose therapy with stem-cell support or combined-modality therapy with chemotherapy

will enhance the CR rate and TTP in this patient population. In addition, trials should be undertaken to evaluate whether the effect of ibritumomab tiuxetan can be enhanced by the addition of cytokines or unlabeled monoclonal antibodies, since Ansell et al<sup>23</sup> recently demonstrated a 69% ORR in a phase I trial of rituximab and interleukin-12.

Initial trials of single-dose anti-CD20 radioimmunconjugates have clearly demonstrated efficacy in producing tumor responses. In addition, high scores have been obtained in measures of patient satisfaction with treatment and for quality of life. The current challenge is to optimize the integration of ibritumomab tiuxetan radioimmunotherapy into the treatment paradigm of B-cell NHL to maximize both tumor response and patient safety.

# ACKNOWLEDGMENT

The acknowledgment is available online at [www.jco.org](http://www.jco.org).

# REFERENCES

1. McLaughlin P, Grillo-López AJ, Link BK, et al: Rituximab chimeric anti-CD20 monoclonal antibody therapy for relapsed indolent lymphoma: Half of patients respond to a four-dose treatment program. *J Clin Oncol* 16:2825-2833, 1998
2. Davis TA, Grillo-López AJ, White CA, et al: Rituximab anti-CD20 monoclonal antibody therapy in non-Hodgkin's lymphoma: Safety and efficacy of re-treatment. *J Clin Oncol* 18:3135-3143, 2000
3. Coiffier B, Lepage E, Herbst R, et al: MabThera (rituximab) plus CHOP is superior to CHOP alone in elderly patients with diffuse large B-cell lymphomas (DLCL): Interim results of a randomized GELA trial. *Blood* 96:223a, 2000 (abstr 550)
4. Czuczman MS: CHOP plus rituximab chemimmunotherapy of indolent B-cell lymphoma. *Semin Oncol* 26:88-96, 1999
5. Czuczman MS, Grillo-López AJ, White CA, et al: Treatment of patients with low-grade B-cell lymphoma with the combination of chimeric anti-CD20 monoclonal antibody and CHOP chemotherapy. *J Clin Oncol* 17:268-276, 1999
6. Vose JM, Link BK, Grossbard ML, et al: Phase II study of rituximab in combination with CHOP chemotherapy in patients with previously untreated, aggressive non-Hodgkin's lymphoma. *J Clin Oncol* 19:389-397, 2001
7. Harris N, Jaffe E, Stein H, et al: A revised European-American classification of lymphoid neoplasms: A proposal from the International Lymphoma Study Group. *Blood* 84:1361-1392, 1994
8. Witzig TE, Gordon LI, Cabanillas F, et al: Randomized controlled trial of yttrium-90-labeled ibritumomab tiuxetan radioimmunotherapy versus rituximab immunotherapy for patients with relapsed or refractory low-grade, follicular, or transformed B-cell non-Hodgkin's lymphoma. *J Clin Oncol* 20:2453-2463, 2002
9. Witzig TE, White CA, Wiseman GA, et al: Phase I/II trial of IDEC-Y2B8 radioimmunotherapy for treatment of relapsed or refractory CD20+ B-cell non-Hodgkin's lymphoma. *J Clin Oncol* 17:3793-3803, 1999
10. Chinn PC, Leonard JE, Rosenberg J, et al: Preclinical evaluation of <sup>90</sup>Y-labeled anti-CD20 monoclonal antibody for treatment of non-Hodgkin's lymphoma. *Int J Oncol* 15:1017-1025, 1999
11. Chinn PC, Morena RA, Wiseman GA, et al: Clinical experience with preparation of Zevalin (ibritumomab tiuxetan, IDEC-Y2B8/IDEC-In2B8) using a radiolabeled kit. *Eur J Nucl Med* 27:1163, 2000 (abstr)
12. Cheson BD, Horning SJ, Coiffier B, et al: Report of an international workshop to standardize response criteria for non-Hodgkin's lymphoma. *J Clin Oncol* 17:1244-1253, 1999
13. Kaplan E, Meier P: Nonparametric estimation for incomplete observations. *J Am Stat Assoc* 53:456-481, 1958
14. Cella D, Tulsky D, Gray G, et al: The Functional Assessment of Cancer Therapy scale: Development and validation of the general measure. *J Clin Oncol* 11:570-579, 1993
15. Wiseman GA, Leigh B, Erwin WD, et al: Radiation dosimetry results for Zevalin radioimmunotherapy of rituximab-refractory non-Hodgkin lymphoma. *Cancer* 94:1349-1357, 2001 (4 suppl)
16. Johnson P, Rohatiner A, Whelan J, et al: Patterns of survival in patients with recurrent follicular lymphoma: A 20-year study from a single center. *J Clin Oncol* 13:140-147, 1995
17. Bastion Y, Sebbon C, Berger F, et al: Incidence, predictive factors, and outcome of lymphoma transformation in follicular lymphoma patients. *J Clin Oncol* 15:1587-1594, 1997
18. Gallagher C, Gregory W, Jones A, et al: Follicular lymphoma: Prognostic factors for response and survival. *J Clin Oncol* 4:1470-1480, 1986
19. Greene MH, Young RC, Merrill JM, et al: Evidence of a treatment dose response in acute nonlymphocytic leukemias which occur after therapy of non-Hodgkin's lymphoma. *Cancer Res* 43:1891-1898, 1983
20. Pedersen-Bjergaard J, Ersboll J, Sorensen HM, et al: Risk of acute nonlymphocytic leukemia and preleukemia in patients treated with cyclophosphamide for non-Hodgkin's lymphomas: Comparison with results obtained in patients treated for Hodgkin's disease and ovarian carcinoma with other alkylating agents. *Ann Intern Med* 103:195-200, 1985
21. Travis LB, Curtis RE, Boice JD Jr, et al: Second cancers following non-Hodgkin's lymphoma. *Cancer* 67:2002-2009, 1991

RADIOIMMUNOTHERAPY FOR RITUXIMAB-REFRACTORY NHL

3269

22. Travis L, Curtis R, Glimelius B, et al: Bladder and kidney cancer following cyclophosphamide therapy for non-Hodgkin's lymphoma. *J Natl Cancer Inst* 87:524-530, 1995
23. Travis LB, Weeks J, Curtis RE, et al: Leukemia following low-dose total body irradiation and chemotherapy for non-Hodgkin's lymphoma. *J Clin Oncol* 14:565-571, 1996
24. Dillman RO, Beauregard JC, Halpern SE, et al: Toxicities and side effects associated with intravenous infusions of murine monoclonal antibodies. *J Biol Response Mod* 5:73-84, 1986
25. Khazaeli M, Saleh M, Wheeler R, et al: Phase I trial of multiple large doses of murine monoclonal antibody CO17-1A. II. Pharmacokinetics and immune response. *J Natl Cancer Inst* 80:937-942, 1988
26. Meeker T, Lowder J, Maloney D, et al: A clinical trial of anti-idiotypic therapy for B cell malignancy. *Blood* 65:1349-1363, 1985
27. Knox SJ, Goris ML, Trisler K, et al: Yttrium-90-labeled anti-CD20 monoclonal antibody therapy of recurrent B-cell lymphoma. *Clin Cancer Res* 2:457-470, 1996
28. Witzig TE, White CA, Gordon L, et al: Response to Zevalin is superior to response to rituximab regardless of age and extent of disease. *Proc Am Soc Clin Oncol* 20:279a, 2001 (abstr 1115)
29. Witzig TE, Gordon LI, Wiseman GA, et al: Reduced dose Zevalin is safe and effective in patients with relapsed or refractory, low grade, follicular or CD20+ transformed B cell non-Hodgkin's lymphoma (L/F/T NHL) and mild thrombocytopenia. *Blood* 96:728a, 2000 (abstr 3149)
30. Wiseman GA, White CA, Stabin M, et al: Phase III <sup>90</sup>Y Zevalin<sup>TM</sup> (yttrium-90 ibritumomab tiuxetan, IDEC-Y2B8) radioimmunotherapy dosimetry results in relapsed or refractory non-Hodgkin's lymphoma. *Eur J Nucl Med* 27:766-777, 2000
31. Wiseman GA, White CA, Sparks RB, et al: Biodistribution and dosimetry results from a phase III prospectively randomized controlled trial of Zevalin radioimmunotherapy for low-grade, follicular, or transformed B-cell non-Hodgkin's lymphoma. *Crit Rev Oncol Hematol* 39:181-194, 2001
32. Witzig TE, Gordon LI, Leigh BR, et al: Baseline peripheral blood B-cell levels correlate with disease-free interval and response to Zevalin radioimmunotherapy of relapsed non-Hodgkin's lymphoma. *Blood* 96:252b, 2000 (abstr 4818)
33. Ansell SM, Witzig TE, Kurtin PJ, et al: Phase I study of interleukin-12 in combination with rituximab in patients with B-cell non-Hodgkin lymphoma. *Blood* 99:67-74, 2002

# Journal of Clinical Oncology

The Official Journal of the American Society of Clinical Oncology

Vol 20, No 15

CONTENTS

August 1, 2002

## EDITORIALS

- Intravesical Therapy for Superficial Cancer: Need for More Options** ..... E. David Crawford 3185
- Challenge of Locally Persistent Prostate Cancer: An Unresolved Clinical Dilemma**  
Christopher J. Logothetis 3187
- Predicting Prostate-Specific Antigen Recurrence Established: Now, Who Will Survive?**  
Anthony V. D'Amico 3188
- Bisphosphonates and Tumor Burden** ..... Gregory R. Mundy 3191

## ORIGINAL REPORTS

### Genitourinary Cancer

- Phase I Trial of Intravesical Gemcitabine in Bacillus Calmette-Guérin-Refractory Transitional-Cell Carcinoma of the Bladder** ..... Guido Dalbagni, Paul Russo, Joel Sheinfeld, Madhu Mazumdar, William Tong, Farhang Rabbani, Machel S. Donat, Harry W. Herr, Pramod Sogani, Dominique dePalma, and Dean Bajorin 3193
- Radical Radiation for Localized Prostate Cancer: Local Persistence of Disease Results in a Late Wave of Metastases** ..... John J. Coen, Anthony L. Zietman, Harjot Thakral, and William U. Shipley 3199
- International Validation of a Preoperative Nomogram for Prostate Cancer Recurrence After Radical Prostatectomy** ..... Markus Graefen, Pierre I. Karakiewicz, Ilias Cagiannos, David I. Quinn, Susan M. Henshall, John J. Grygiel, Robert L. Suherland, Phillip D. Stricker, Eric Klein, Patrick Kupelian, Donald G. Skinner, Gary Lieskovsky, Bernard Bochner, Hartwig Huland, Peter G. Hammerer, Alexander Haese, Andreas Erbersdobler, James A. Eastham, Jean de Kernion, Thomas Cangiano, Fritz H. Schröder, Mark F. Wildhagen, Theo H. van der Kwast, Peter T. Scardino, and Michael W. Kattan 3206

*Journal of Clinical Oncology* (ISSN 0732-183X) is published 24 times a year, twice monthly, by Lippincott Williams & Wilkins, 351 West Camden Street, Baltimore, MD 21201-2436. Periodicals postage paid at Hagerstown, MD, and at additional mailing offices. The GST number for Canadian Subscribers is 895524239.

Editorial correspondence should be addressed to Daniel G. Haller, MD, *Journal of Clinical Oncology*, 330 John Carlyle St, Suite 300, Alexandria, VA 22314. Telephone: (703) 797-1900; FAX: (703) 684-8720. Email: jco@asco.org. Internet: <http://www.jco.org>

**POSTMASTER:** ASCO members send change of address to American Society of Clinical Oncology, 1900 Duke St, Suite 200, Alexandria, VA 22314. Non-members send change of address to *Journal of Clinical Oncology*, c/o Lippincott Williams & Wilkins, PO Box 1550, Hagerstown, MD 21740.

Yearly subscription rates: United States and possessions: individual, \$325.00; institution, \$452.00; single issue, \$25.00. Canada and Mexico: individual, \$456.00; institution, \$583.00; single issue, \$25.00. All other countries: individual, \$496.00; institution, \$623.00; single issue, \$25.00. Student and resident: United States and possessions: \$127.00, Canada and Mexico: \$153.00, all other countries: \$193.00. To receive student/resident rate, orders must be accompanied by name of affiliated institution, date of term, and the signature of program/residency coordinator on institution letterhead. Orders will be billed at individual rate until proof of status is received. Current prices are in effect for back volumes and back issues. Back issues sold in conjunction with a subscription rate are on a prorated basis. Subscriptions are accepted on a calendar year basis. Prices are subject to change without notice. Single issues, both current and back, exist in limited quantities and are offered for sale subject to availability. Publication Mail Agreement Number 363289.

1

2

**Rx Only**

3

**BEXXAR®**

4

**Tositumomab and Iodine I 131 Tositumomab**

5

**WARNINGS**

6 **Hypersensitivity Reactions, including Anaphylaxis:** Medications for the  
7 treatment of severe hypersensitivity reactions should be available for immediate  
8 use. Patients who develop severe hypersensitivity reactions should have  
9 infusions of the BEXXAR therapeutic regimen discontinued and receive medical  
10 attention (See **WARNINGS**).

11 **Prolonged and Severe Cytopenias:** The majority of patients who received the  
12 BEXXAR therapeutic regimen experienced severe thrombocytopenia and  
13 neutropenia. The BEXXAR therapeutic regimen should not be administered to  
14 patients with >25% lymphoma marrow involvement and/or impaired bone marrow  
15 reserve (See **WARNINGS** and **ADVERSE REACTIONS**).

16 **Pregnancy Category X:** The BEXXAR therapeutic regimen can cause fetal  
17 harm when administered to a pregnant woman.

18 **Special requirements:** The BEXXAR therapeutic regimen (Tositumomab and  
19 Iodine I 131 Tositumomab) contains a radioactive component and should be  
20 administered only by physicians and other health care professionals qualified by  
21 training in the safe use and handling of therapeutic radionuclides. The BEXXAR  
22 therapeutic regimen should be administered only by physicians who are in the  
23 process of being or have been certified by Corixa Corporation in dose calculation  
24 and administration of the BEXXAR therapeutic regimen.

25

26

**DESCRIPTION**

27

The BEXXAR therapeutic regimen (Tositumomab and Iodine I 131

28

Tositumomab) is an anti-neoplastic radioimmunotherapeutic monoclonal

29 antibody-based regimen composed of the monoclonal antibody,  
30 Tositumomab, and the radiolabeled monoclonal antibody, Iodine I 131  
31 Tositumomab.

## 32 **Tositumomab**


33 Tositumomab is a murine IgG<sub>2a</sub> lambda monoclonal antibody directed  
34 against the CD20 antigen, which is found on the surface of normal and  
35 malignant B lymphocytes. Tositumomab is produced in an antibiotic-free  
36 culture of mammalian cells and is composed of two murine gamma 2a  
37 heavy chains of 451 amino acids each and two lambda light chains of 220  
38 amino acids each. The approximate molecular weight of Tositumomab is  
39 150 kD.

40 Tositumomab is supplied as a sterile, pyrogen-free, clear to opalescent,  
41 colorless to slightly yellow, preservative-free liquid concentrate. It is  
42 supplied at a nominal concentration of 14 mg/mL Tositumomab in 35 mg  
43 and 225 mg single-use vials. The formulation contains 10% (w/v) maltose,  
44 145 mM sodium chloride, 10 mM phosphate, and Water for Injection, USP.  
45 The pH is approximately 7.2.

## 46 **Iodine I 131 Tositumomab**

47 Iodine I 131 Tositumomab is a radio-iodinated derivative of Tositumomab  
48 that has been covalently linked to Iodine-131. Unbound radio-iodine and  
49 other reactants have been removed by chromatographic purification steps.  
50 Iodine I 131 Tositumomab is supplied as a sterile, clear, preservative-free  
51 liquid for IV administration. The dosimetric dosage form is supplied at  
52 nominal protein and activity concentrations of 0.1 mg/mL and 0.61 mCi/mL  
53 (at date of calibration), respectively. The therapeutic dosage form is  
54 supplied at nominal protein and activity concentrations of 1.1 mg/mL and  
55 5.6 mCi/mL (at date of calibration), respectively. The formulation for the  
56 dosimetric and the therapeutic dosage forms contains 5.0%–6.0% (w/v)  
57 povidone, 1–2 mg/mL maltose (dosimetric dose) or 9–15 mg/mL maltose  
58 (therapeutic dose), 0.85–0.95 mg/mL sodium chloride, and 0.9–1.3 mg/mL  
59 ascorbic acid. The pH is approximately 7.0.

60



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Br

Search PubMed for Go Clear

Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

1: Int J Oncol. 1999 Nov;15(5):1017-25.

Related Articles, Link

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

**Preclinical evaluation of 90Y-labeled anti-CD20 monoclonal antibody for treatment of non-Hodgkin's lymphoma.****Chinn PC, Leonard JE, Rosenberg J, Hanna N, Anderson DR.**

IDEC Pharmaceuticals Inc., San Diego, CA 92121, USA.

A high-affinity IgG1 kappa murine monoclonal anti-CD20 antibody (IDEC-2B8) was developed for radioimmunotherapy of non-Hodgkin's B-cell lymphoma. A stable immunoconjugate (Zevalintrade mark) was prepared by reacting IDEC-2B8 with a derivative of diethylenetriaminepentaacetic acid, designated MX-DTPA, a chelator exhibiting high affinity and retention for 90Y. Zevalin exhibited antigen specificity, human tissue reactivity, and preclinical safety profile comparable to the native antibody. The conjugate radiolabeled with 90Y (90Y-Zevalin) or 111In (111In-Zevalin) exhibited excellent retention of immunoreactivity with radioincorporations >95%. The radiolabeled conjugates formulated in PBS containing human serum albumin were stable in vitro at 4 degrees C for 48 h as indicated by negligible loss of radioisotope and retention of binding to CD20+ cells. In vitro human serum stability studies at 37 degrees C with 90Y-Zevalin indicated that loss of 90Y from the conjugate was minimal, averaging 1% per day. Biodistribution studies in BALB/c mice confirmed the in vitro stability of 90Y-Zevalin and 111In-Zevalin. In particular, excellent in vivo retention of 90Y by the conjugate was demonstrated by minimal bone accumulation (<=3% of the injected dose over three days). Radiation dose estimates to normal organs calculated from mouse biodistribution studies with 90Y-Zevalin were comparable to those determined in a phase I/II clinical trial and below generally accepted safe radiation levels. Studies in athymic mice bearing CD20+ tumors demonstrated that 111In-Zevalin accumulated in the tumors preferentially compared with normal organs. 90Y-Zevalin is currently being evaluated in phase III clinical trials for treatment of relapsed or refractory low-grade, follicular or transformed B-cell non-Hodgkin's lymphoma.

PMID: 10536187 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text